arXiv:1512.06756v1 [gr-qc] 18 Dec 2015

# Stationary Point of the Hilbert Action Principle

James W. York

Department of Physics
North Carolina State University
Raleigh, NC, 27695-8202

November 11th, 2015

**Abstract**

The Hilbert action principle for the Einstein equations has two boundary terms that must be subtracted in prder to obtain a true stationary point. In this paper, I obtain both of them.

Assume that spacetime $V$ is topologically $\Sigma \times \mathbb{R}$ where $\Sigma$ is a spacelike Cauchy slice. The splittings of the spacetime curvature tensor $R_{\mu\nu\alpha\beta}$ works as follows[1].

$$R_{ijkl} = \bar{R}_{ijkl} + \left(K_{ik}K_{jl} - K_{il}K_{jk}\right), \qquad \text{Gauss} \tag{1}$$

$$R_{0ijk} = N\left(\nabla_j K_{ik} - \nabla_k K_{ij}\right), \qquad \text{Codazzi} \tag{2}$$

$$R_{0i0j} = N\left(\hat{\partial}_0 K_{ij} + K_{ik}K^k_j + \nabla_i\nabla_j N N\right), \qquad \text{Ricci} \tag{3}$$

where $\hat{\partial}_0 = \partial_t - \mathcal{L}_\beta$. $N$ is the lapse function and $\beta^i$ is the shift vector. In (1) $\bar{R}_{ijkl}$ is the Reimann tensor of $\Sigma$ and $\nabla_i$ is the spatial covariant derivative. In three dimmensions $\bar{R}_{ijkl}$ is equivalent to the Ricci Tensor $\bar{R}_{ij}$. $K_{ij}$ is the extrinsic curvature of $\Sigma$. Proceeding with the splitting of spacetime curvatures tensors, I find

$$Ric_{\beta\gamma} = R_{\alpha\beta}{}^{\alpha}{}_{\gamma} \tag{4}$$

$$Ric_{ij} = \bar{R}_{ij} - N^{-1}\hat{\partial}_0 K_{ij} + KK_{ij} - 2K_{ij}K^k{}_j + N^{-1}\nabla_i\nabla_j N, \tag{5}$$

where $K = g_{ij}K^{ij}$

$$Ric_{00} = N\left(\hat{\partial}_0 K - NK_{ij}K^{ij}\Delta N\right), \tag{6}$$

with $\Delta = g^{ij}\nabla_i\nabla_j$.

The spacetime scalar curvature is $R = g^{\alpha\beta}R_{ic\alpha\beta}$ from which

$$R = N\hat{\partial}_0 K - 2N^{-1}\Delta N + \left(\bar{R} + K_{ij}K^{ij} - K^2\right) \tag{7}$$

and

$$\begin{aligned} N\sqrt{g}R &= N\sqrt{g}\left(\bar{R} + K_{ij}K^{ij} - K^2\right) - \partial_t\left(\sqrt{g}K\right) \\ &+ \partial_i\left[\sqrt{g}\left(K\beta^i - g^{ij}\partial_j N\right)\right]. \end{aligned} \tag{8}$$

The tensor $\left(-\partial_t\sqrt{g}k\right)$ was recognized in [2]. The spatial scalar curvature is

$$\bar{R} = R_{(2)} - \left(\chi_{ij}\chi^{ij} + \chi^2\right) + 2\left(\mathcal{L}_s\chi - \alpha^{-1}\bar{g}^{ij}\bar{\nabla}_i\bar{\nabla}_j\alpha\right), \tag{9}$$

where $R_{(2)}$ is the scalar curvature of the reference this space $S$ embedded in $\Sigma$. The metric of $S$ is $\bar{g}_{ij}$ and its covariant derivative is $\bar{\nabla}_i$. Assume that $\Sigma$ is foliated by $S$'s, all of which are outside of any sources. The analogue of $N$ is $\alpha$, a lapse function associated with the foliation of $\Sigma$ by $S$'s. The $S$'s are topologically two-spheres that approach a genuine two-sphere as $r \to \infty$ in asymptotically Euclidean $\Sigma$'s (AES's). Define a unit vector $s^i$ that is the outward-pointing unit normal of the $S$'s. The extensive curvature of $S$ in $\Sigma$ is $\chi_{ij}$; it satisfies

$$\chi_{ij} = (2\alpha)^{-1}\left[\partial_r\bar{g}_{ij} + \left(\bar{\nabla}_i s_j + \bar{\nabla}_j s_i\right)\right]. \tag{10}$$

$\mathcal{L}_N\chi = \mathcal{L}_N\left(\bar{g}^{ij}\chi_{ij}\right)$ has support on AES's because the $S$'s approach a genuine two-sphere in the limit as $r \to \infty$. At asymptotia, the exact form of the Hilbert action principle (with $k = 4\pi G$, $G = 6.670(4) \times 10^{-8}\mathrm{gm}^{-1}\mathrm{cm}^3\mathrm{sec}^{-2}$ and $c = 2.9979250(10) \times 10^{10}\mathrm{cm\ sec}^{-1}$) is

$$\begin{aligned} \left(c^4/4k\right)\int N\sqrt{g}Rd^4k = (c^4/4k)\int[N\sqrt{g}\left(\bar{R} + K_{ij}K^{ij} - K^2\right) \\ +\partial_i\left(\sqrt{g}K\beta^i - g^{ij}\partial_j N\right) - \partial_t\left(\sqrt{g}K\right) - \mathcal{L}_N\chi]d^4\chi. \end{aligned} \tag{11}$$

The other tensors in (9) have no support as $r \to \infty$. Variation of (11) with respect to $g_{\mu\nu}$ gives as a genuine stationary point

$$\left(c^4/4k\right)Ein^{\mu\nu} = \left(c^4/4k\right)\left[Ric^{\mu\nu} - \tfrac{1}{2}g^{\mu\nu}R\right] = 0. \tag{12}$$

$D_\nu Ein^{\mu\nu} \equiv 0$, where $D_\nu$ is the spacetime covariant derivative. Equation (12) gives the vacuum Einstein equations. If sources are present, a term must be added to (11) to produce stress-energy-momentum tensor $T^{\mu\nu}$ that necessarily satisfies $D_\nu T^{\mu\nu} =$. This is not a true conservation law except in the presence of one or more Killing vectors $\xi^\mu$ that satisfy

$$\mathcal{L}_\xi g_{\mu\nu} = D_\nu \xi_\mu + D_\mu \xi_\nu = 0, \tag{13}$$

in which case

$$D_\mu \left( T^{\mu\nu} \xi_\nu \right) = 0 \tag{14}$$

is a true conservation law.

# Acknowledgment

The author thanks J. David Brown for finding a missing page of his CV.

# References

[1] J.A. Schouten, *Ricci-Calculus, second edition (1954), Springer-Verlag*, Berlin-Göttingen-Heidelberg.

[2] J.W. York, Boundary terms in the action principles of general relativity (in honor of the seventy-fifth birthday of John A. Wheeler). Foundations of Physics 16:249-258 (1986).